

The eigenvalues of q -Kneser graphs

Benjian Lv Kaishun Wang*

Sch. Math. Sci. & Lab. Math. Com. Sys., Beijing Normal University, Beijing, 100875, China

Abstract

In this note, we prove some combinatorial identities and obtain a simple form of the eigenvalues of q -Kneser graphs.

AMS classification: 05A19, 05E30

Key words: q -Kneser graph; eigenvalue.

Let q be a prime power. For any integer n and positive integer i , the Gaussian coefficients is defined by

$$\begin{bmatrix} n \\ i \end{bmatrix}_q = \prod_{j=0}^{i-1} \frac{q^{n-j} - 1}{q^{i-j} - 1}.$$

By convention $\begin{bmatrix} n \\ 0 \end{bmatrix}_q = 1$ for every integer n . From now on, we will omit the subscript q . Note that

$$\begin{bmatrix} n \\ i \end{bmatrix} = \begin{bmatrix} n-1 \\ i-1 \end{bmatrix} + q^i \begin{bmatrix} n-1 \\ i \end{bmatrix}. \quad (1)$$

Let $\mathbb{F}_q^v$ be a v -dimensional vector space over a finite field $\mathbb{F}_q$. The q -Kneser graph $qK(v, k)$ has as vertex set the collection of k -dimensional subspaces of $\mathbb{F}_q^v$. Two vertices are adjacent if they intersect trivially. If $k \leq v < 2k$, then $qK(v, k)$ is null graph, so we only consider the case $v \geq 2k$.

Delsarte [1] calculated the eigenvalues of Grassmann schemes. In particular, the eigenvalues of $qK(v, k)$ were given.

Theorem 1 ([1, Theorem 10]) *All the distinct eigenvalues of $qK(v, k)$ are*

$$\lambda_j = (-1)^j q^{(k-j)j + \binom{j}{2}} \sum_{s=0}^{k-j} (-1)^s q^{\binom{s}{2}} \begin{bmatrix} k-j \\ s \end{bmatrix} \begin{bmatrix} v-2j-s \\ v-k-j \end{bmatrix}, \quad (2)$$

where $j = 0, 1, \dots, k$.

*Corresponding author. E-mail address: wangks@bnu.edu.cn

The eigenvalues of Kneser graphs are deduced to a simple form in [2, Theorem 9.4.3]. In this note, we shall generalize this result to vector spaces, and obtain a simple form of eigenvalues of $qK(v, k)$. We start with some useful combinatorial identities.

Lemma 2 *For any integer n and nonnegative integer i , we have*

$$\begin{bmatrix} n \\ i \end{bmatrix} = (-1)^i q^{ni - \binom{i}{2}} \begin{bmatrix} -n + i - 1 \\ i \end{bmatrix}.$$

Proof. If $i = 0$, the identity is obvious. If $i > 0$, then

$$\begin{aligned} \begin{bmatrix} n \\ i \end{bmatrix} &= (-1)^i \prod_{j=0}^{i-1} \frac{1 - q^{n-j}}{q^{i-j} - 1} \\ &= (-1)^i q^{\frac{(2n-i+1)i}{2}} \prod_{j=0}^{i-1} \frac{q^{-n+j} - 1}{q^{i-j} - 1} \\ &= (-1)^i q^{ni - \binom{i}{2}} \prod_{j=0}^{i-1} \frac{q^{-n+i-1-j} - 1}{q^{i-j} - 1} \\ &= (-1)^i q^{ni - \binom{i}{2}} \begin{bmatrix} -n + i - 1 \\ i \end{bmatrix}, \end{aligned}$$

as desired. □

The following identity is a generalization of [3, Theorem 2.14].

Lemma 3 *For any integer n and nonnegative integer a , we have*

$$\sum_{s=0}^a (-1)^s q^{\binom{s}{2}} \begin{bmatrix} n \\ s \end{bmatrix} = q^{na} \begin{bmatrix} a - n \\ a \end{bmatrix}.$$

Proof. We prove the result by induction on a . If $a = 0$, then the result is trivial. Suppose $a \geq 1$. By induction and Lemma 2,

$$\begin{aligned} \sum_{s=0}^a (-1)^s q^{\binom{s}{2}} \begin{bmatrix} n \\ s \end{bmatrix} &= \sum_{s=0}^{a-1} (-1)^s q^{\binom{s}{2}} \begin{bmatrix} n \\ s \end{bmatrix} + (-1)^a q^{\binom{a}{2}} \begin{bmatrix} n \\ a \end{bmatrix} \\ &= q^{n(a-1)} \begin{bmatrix} a - 1 - n \\ a - 1 \end{bmatrix} + q^{na} \begin{bmatrix} a - 1 - n \\ a \end{bmatrix} \\ &= q^{na} \begin{bmatrix} a - n \\ a \end{bmatrix}. \end{aligned}$$

Hence, the desired result follows. □

Lemma 4 Let m, a, t be nonnegative integers with $t \leq a \leq m$. Then

$$\sum_{s=0}^a (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-s \\ t \end{bmatrix} = q^{m(a-t)} \begin{bmatrix} a-m \\ a-t \end{bmatrix}. \quad (3)$$

Proof. We prove the result by induction on a and t . If $t = 0$, (3) is immediate by Lemma 3. If $a = t$, (3) is straightforward. Now suppose $1 \leq t < a$. By (1) and induction,

$$\begin{aligned} & \sum_{s=0}^a (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-s \\ t \end{bmatrix} \\ &= \sum_{s=0}^{a-1} (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-s \\ t \end{bmatrix} \\ &= \sum_{s=0}^{a-1} (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-1-s \\ t-1 \end{bmatrix} + q^t \sum_{s=0}^{a-1} (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-1-s \\ t \end{bmatrix} \\ &= q^{m(a-t)} \begin{bmatrix} a-1-m \\ a-t \end{bmatrix} + q^t q^{m(a-1-t)} \begin{bmatrix} a-1-m \\ a-1-t \end{bmatrix} \\ &= q^{m(a-t)} \begin{bmatrix} a-m \\ a-t \end{bmatrix}. \end{aligned}$$

Hence, the desired result follows. $\square$

Theorem 5 Let m, a, t be nonnegative integers with $a \geq m$ and $a \geq t$. Then

$$\sum_{s=0}^m (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-s \\ t \end{bmatrix} = q^{m(a-t)} \begin{bmatrix} a-m \\ a-t \end{bmatrix}. \quad (4)$$

Proof. We prove the result by induction on a and t . If $t = 0$, (4) is immediate from Lemma 3. If $a = m$, (4) holds by Lemma 4. If $a = t$, (4) is straightforward. Suppose $a \geq m + 1$ and $1 \leq t < a$. By (1) and induction,

$$\begin{aligned} & \sum_{s=0}^m (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-s \\ t \end{bmatrix} \\ &= \sum_{s=0}^m (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-1-s \\ t-1 \end{bmatrix} + q^t \sum_{s=0}^m (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-1-s \\ t \end{bmatrix} \\ &= q^{m(a-t)} \begin{bmatrix} a-1-m \\ a-t \end{bmatrix} + q^t q^{m(a-1-t)} \begin{bmatrix} a-1-m \\ a-t-1 \end{bmatrix} \\ &= q^{m(a-t)} \begin{bmatrix} a-m \\ a-t \end{bmatrix}. \end{aligned}$$

Therefore, (4) holds. $\square$

Substituting $t = a - m$ in (4), we obtain

Corollary 6 For nonnegative integers $a \geq m$, we have

$$\sum_{s=0}^m (-1)^s q^{\binom{s}{2}} \begin{bmatrix} m \\ s \end{bmatrix} \begin{bmatrix} a-s \\ a-m \end{bmatrix} = q^{m^2} \begin{bmatrix} a-m \\ m \end{bmatrix}.$$

Next, we shall deduce a simple form of the eigenvalues λ_j in (2) of $qK(v, k)$.

Theorem 7 All the distinct eigenvalues of $qK(v, k)$ are

$$\lambda_j = (-1)^j q^{\binom{k}{2} + \binom{k-j+1}{2}} \begin{bmatrix} v-k-j \\ v-2k \end{bmatrix}, \quad j = 0, 1, \dots, k.$$

Moreover, the multiplicity of λ_j is 1 if $j = 0$, and $\begin{bmatrix} v \\ j \end{bmatrix} - \begin{bmatrix} v \\ j-1 \end{bmatrix}$ if $j \geq 1$.

Proof. By (2) and Corollary 6, we have

$$\begin{aligned} \lambda_j &= (-1)^j q^{\binom{k-j}{2} + \binom{j}{2}} \sum_{s=0}^{k-j} (-1)^s q^{\binom{s}{2}} \begin{bmatrix} k-j \\ s \end{bmatrix} \begin{bmatrix} v-2j-s \\ v-2j-(k-j) \end{bmatrix} \\ &= (-1)^j q^{\binom{k-j}{2} + \binom{j}{2}} q^{\binom{k-j}{2}} \begin{bmatrix} v-2j-(k-j) \\ k-j \end{bmatrix} \\ &= (-1)^j q^{\binom{k}{2} + \binom{k-j+1}{2}} \begin{bmatrix} v-k-j \\ v-2k \end{bmatrix}. \end{aligned}$$

By similar arguments in [1, 4], the multiplicity of each λ_j may be computed. $\square$

Acknowledgement

This research is partially supported by NCET-08-0052, NSF of China (10871027) and the Fundamental Research Funds for the Central Universities of China.

References

- [1] P. Delsarte, Association schemes and t -designs in regular semilattices, J. Combin. Theory Ser. A, 20 (1976) 230-243.
- [2] C.D. Godsil and G. Royle, Algebraic Graph Theory, Springer-Verlag, New York, Berlin, Heidelberg, 2001.
- [3] Z. Wan, Binomial coefficient and Gaussian binomial coefficient, Mathematics Bulletin, 11 (1994) 7-13 (in chinese).
- [4] R.M. Wilson, The exact bound in the Erdős-Ko-Rado theorem, Combinatorica, 4 (1984) 247-257.